



E1630
JACC March 12, 2013
Volume 61, Issue 10

TCT@ACC-i2: Invasive and Interventional Cardiology

THE ROLE OF FEMALE SEX IN THE CONTEMPORARY TREATMENT OF THE LEFT MAIN CORONARY ARTERY INSIGHTS FROM THE W- DELTA (WOMEN- DRUG ELUTING STENT FOR LEFT MAIN CORONARY ARTERY DISEASE) REGISTRY

Oral Contributions

West, Room 2022

Saturday, March 09, 2013, 8:45 a.m.-8:55 a.m.

Session Title: Left Main and Bifurcation PCI

Abstract Category: 46. TCT@ACC-i2: Coronary Intervention, LM/Bifurcations

Presentation Number: 2901-4

Authors: *Gill Louise Buchanan, Alaide Chieffo, Emanuele Meliga, Seung-Jung Park, Yoshinobu Onuma, Piera Capranzano, Marco Valgimigli, Sanda Jegere, Raj Makkar, Igor Palacios, Pawel Buszman, Imad Sheiban, Roxana Mehran, Christoph Naber, Ronan Margey, Arvind Agnihotri, Davide Capodanno, Martin Leon, Jean Fajadet, Marie Morice, Thierry Lefevre, Andrejs Erglis, Corrado Tamburino, Patrick Serruys, Antonio Colombo, San Raffaele Scientific Institute, Milan, Italy*

Background: Limited data are available on the optimal treatment of women with unprotected left main coronary artery (ULMCA) disease. The aim was to specifically evaluate long-term clinical outcomes in women treated with percutaneous coronary intervention (PCI) with drug-eluting stents (DES) vs. coronary artery bypass grafting (CABG).

Methods: All consecutive women from the Drug Eluting stent for Left main coronary Artery disease (DELTA) Registry with ULMCA disease treated by PCI with DES or CABG were analyzed. A propensity matching was performed to adjust for baseline differences between the 2 treatment groups.

Results: In total, 818 women were included: 489 (59.8%) underwent treatment with PCI with DES vs. 329 (40.2%) with CABG. Propensity score matching identified 176 matched pairs of patients, with no differences in the primary study objective of death, myocardial infarction (MI) or cerebrovascular accident (CVA) (Hazard Ratio [HR] 0.711; 95% Confidence Interval [CI] 0.387-1.308; $p=0.273$), all-cause (HR 0.722; 95% CI 0.357-1.461; $p=0.365$) or cardiovascular mortality (HR 1.100; 95% CI 0.455-2.660; $p=0.832$), MI (HR 0.362; 95% CI 0.094-1.388) or CVA (HR 1.200; 95% CI 0.359-4.007; $p=0.767$). However, there was an advantage of CABG over PCI in major adverse cardiovascular and cerebrovascular events (MACCE) (HR 0.429; 95% CI 0.254-0.723; $p=0.001$), driven exclusively by target vessel revascularization (TVR) (HR 0.185; 95% CI 0.079-0.432; $p<0.001$).

Conclusions: In women with significant ULMCA disease, no difference was observed in death, MI and stroke at long term follow-up. However, repeat revascularization remained higher in the PCI cohort.